

LOCAL-GLOBAL PRINCIPLE FOR THE BAUM-CONNES CONJECTURE WITH COEFFICIENTS

PAUL BAUM, STEPHEN MILLINGTON AND ROGER PLYMEN

ABSTRACT. We establish the Hasse principle (local-global principle) in the context of the Baum-Connes conjecture with coefficients. We illustrate this principle with the discrete group $GL(2, F)$ where F is any global field.

1. INTRODUCTION

Let G be a second countable locally compact Hausdorff topological group. We shall say that G satisfies *BCC*, or *BCC* is true for G , if the Baum-Connes conjecture with coefficients in an arbitrary $G - C^*$ -algebra is true for G .

Our first result is the following permanence property:

Theorem 1.1. *Let G be the ascending union of open subgroups G_n , and let A be a $G - C^*$ -algebra. If each subgroup G_n satisfies the Baum-Connes conjecture with coefficients A , then G satisfies the Baum-Connes conjecture with coefficients A .*

Our main application is to the following new permanence property.

Theorem 1.2. *Let F be a global field, $\mathbb{A}$ its ring of adeles, G a linear algebraic group defined over F . Let F_v denote a place of F . If *BCC* is true for each local group $G(F_v)$ then *BCC* is true for the adelic group $G(\mathbb{A})$.*

Another application of Theorem 1.1 is the proof of the Baum-Connes conjecture for reductive adelic groups [2].

To derive Theorem 1.2 from Theorem 1.1 we note that the adelic group $G(\mathbb{A})$ admits an ascending union of open subgroups. We then make use of a crucial permanence property due to Chabert-Echterhoff [5], namely that *BCC* is stable under direct product of finitely many groups.

If G satisfies *BCC*, then any closed subgroup of G also satisfies *BCC* [5, Theorem 2.5]. Since $G(F)$ is a discrete subgroup of $G(\mathbb{A})$, we have the following result:

Date: version 6: February 1, 2008.

Theorem 1.3. *If BCC is true for each local group $G(F_v)$ then BCC is true for the discrete group $G(F)$.*

There is, at present, a limited supply of local groups for which BCC is known to be true. Nevertheless, some examples are known.

For the group $SO(n, 1)$, Kasparov [12] proved that $\gamma = 1_G$ in the Kasparov representation ring $R(G) = KK_G(\mathbb{C}, \mathbb{C})$. For the group $SU(n, 1)$, Julg-Kasparov [11] and Higson-Kasparov [8] proved that $\gamma = 1_G$ in the ring $R(G)$. For the group $Sp(n, 1)$, Julg [10] has recently proved that, for any $G - C^*$ -algebra A , the image of γ via the map:

$$R(G) \rightarrow KK_G(A, A) \rightarrow KK(A \rtimes_r G, A \rtimes_r G) \rightarrow \text{End } K_*(A \rtimes_r G)$$

is the identity element even though $\gamma \neq 1$ in the ring $R(G)$. Therefore the following rank-one Lie groups satisfy BCC:

$$SO(n, 1), SU(n, 1), Sp(n, 1).$$

Let F be a global field and consider the group $GL(2, F)$. At each place v of F we have the local group $GL(2, F_v)$. The Lie group $SL(2, \mathbb{R})$ acts properly and isometrically on the Poincaré disc, and so has the Haagerup property [9, p.159]. The Lie group $SL(2, \mathbb{C})$ acts properly and isometrically on the Poincaré ball [7, p.11], and so has the Haagerup property [9, p.159]. If v is a finite place then the totally disconnected group $SL(2, F_v)$ acts properly and isometrically on its tree [13], admits a proper 1-cocycle, and so has the Haagerup property [9, p.155].

For each place v , the determinant map creates a short exact sequence

$$1 \rightarrow SL(2, F_v) \rightarrow GL(2, F_v) \rightarrow F_v^\times \rightarrow 1$$

in which $SL(2, F_v)$ is a closed normal subgroup with the Haagerup property, and the locally compact abelian group $F_v^\times$ certainly satisfies BCC. Therefore, by [5, Corollary 3.14] each local group $GL(2, F_v)$ satisfies BCC.

Let $\mathbb{A}_F$ be the adèle ring attached to F . By Theorem 1.2, we have

Theorem 1.4. *The adelic group $GL(2, \mathbb{A}_F)$ satisfies BCC.*

By Theorem 1.3, we have

Theorem 1.5. *The discrete group $GL(2, F)$, and each of its subgroups, satisfies BCC.*

Our method is therefore an example of the Hasse principle (local-global principle) applied to the *local* groups $G(F_v)$ and the *discrete* group $G(F)$.

It is worth noting that if the Baum-Connes conjecture fails for the discrete group $SL(n, \mathbb{Z})$, then BCC fails for $SL(n, \mathbb{R})$.

The discrete group $SL(2, F)$, with F a global field, is discussed in the Bourbaki seminar by Pierre Julg [9, p. 160]. He proves that $SL(2, F)$ has the Haagerup property. It then follows from a theorem of Higson-Kasparov [8] that $SL(2, F)$ satisfies the Baum-Connes conjecture, see [9, p.152].

In the course of our work, we find it necessary to use a model $P_c(G)$ of the universal example for proper actions of G which is itself a direct limit of compact spaces. This model $P_c(G)$ is paracompact, Hausdorff and separable, but not metrizable, and so falls outside the discussion of proper actions in [1]. We have therefore to choose a different starting point for the theory of proper actions: we use the definition of Bourbaki [4].

This paper is a sequel to our Note [2]. We thank Siegfried Echterhoff, Pierre Julg, Ryszard Nest and Richard Sharp for valuable conversations.

2. ADELIC GROUPS

A *local field* is a non discrete locally compact topological field. It is shown in [15] that a local field F must be of the following form. If $\text{char}(F) = 0$, then $F = \mathbb{R}, \mathbb{C}$ or a finite extension of $\mathbb{Q}_p$ for some prime p . If $\text{char}(F) = p > 0$, then F is the field $\mathbb{F}_q((X))$ of formal Laurent series (with finite tail) in one variable with coefficients in a finite field $\mathbb{F}_q$.

The fields $\mathbb{R}, \mathbb{C}$ are known as *archimedean fields*. All other local fields are known as *nonarchimedean fields*. The topology on a nonarchimedean field is always totally disconnected.

Let $\mathbb{F}_p(t)$ denote the field of fractions of the polynomial ring $\mathbb{F}_p[t]$. A *global field* is a finite extension of $\mathbb{Q}$, or a finite extension of the function field $\mathbb{F}_p(t)$. A *completion* (v, F_v) of F is a dense isomorphic embedding v of F into a local field F_v . Two completions $(v, F_v), (u, F_u)$ are said to be equivalent if there is an isomorphism ρ of F_v onto F_u such that $u = \rho \circ v$. A *place* of F is an equivalence class of completions. We say the place (v, F_v) is *infinite* if F_v is an archimedean field and *finite* otherwise. If $\text{char}(F) = p > 0$, then F has countably many finite places and *no* infinite places. If $\text{char}(F) = 0$ then F has countably many finite places and finitely many (but at least one) infinite places.

Suppose we have an ascending sequence of topological spaces

$$X_1 \subset X_2 \subset X_3 \subset \dots$$

Then we can give the union $X = \cup X_n$ the *direct limit topology*: that is a set is open in X if and only if it has open intersection with each X_n .

The following proposition admits a standard proof in point-set topology.

Lemma 2.1. *Let $X_1 \subset X_2 \subset X_3 \subset \dots$ be a sequence of T_1 (=points are closed) topological spaces and give $X = \cup_{n=1}^{\infty} X_n$ the direct limit topology. Then any compact subset of X lies entirely within some X_n .*

Suppose that for each $n = 1, 2, 3, \dots$ we have a locally compact, second countable and Hausdorff topological group G_n , such that G_m is an open subgroup of G_n for $m \leq n$:

$$G_1 \subset G_2 \subset G_3 \subset \dots$$

Let $G = \cup_{n=1}^{\infty} G_n$ and furnish this with the direct limit topology. Then G is a topological group which is locally compact, second countable and Hausdorff.

Any nonarchimedean local field F_v contains a unique maximal compact open subring $\mathcal{O}_v$. Let S denote any finite set of places of F which contains all the infinite places. By an *adele* we mean an element $a = (a)_v$ of the product $\prod_v F_v$ such that $a \in A_S = \prod_{v \in S} F_v \times \prod_{v \notin S} \mathcal{O}_v$ for some S . The adeles of F form a ring $\mathbb{A}_F$, addition and multiplication being defined componentwise. Each A_S has its natural topology and $\mathbb{A}_F = \cup_S A_S$ is topologized as the inductive limit with respect to S . There is an obvious embedding of F in $\mathbb{A}_F$, by means of which we identify F with a subring of $\mathbb{A}_F$. The field F is a discrete cocompact subfield of the non-discrete locally compact semisimple commutative ring $\mathbb{A}_F$.

Now suppose G is a linear algebraic group defined over F . We shall be interested in the *adelic group* $G(\mathbb{A}_F)$ of $\mathbb{A}_F$ -rational points of G . For a finite place v of F let $G(\mathcal{O}_v)$ denote the group $G(F_v) \cap GL_n(\mathcal{O}_v)$. We set

$$G_S = \prod_{v \in S} G(F_v) \times \prod_{v \notin S} G(\mathcal{O}_v)$$

Then $G(\mathbb{A})$ is equal, by definition, to the direct limit of the groups G_S . We now equip $G(\mathbb{A})$ with the direct limit topology, following Weil [14, p.2]. Then $G(\mathbb{A})$ is a locally compact second countable Hausdorff group. The map $x \mapsto (x, x, x, \dots)$ embeds $G(F)$ as a *discrete* subgroup of $G(\mathbb{A})$.

3. PROPER ACTIONS AND UNIVERSAL EXAMPLES

We recall that a topological space X is *completely regular* if it satisfies the following separation axiom: If B is a closed subset of X and $p \in$

$X \setminus B$ then there exists a continuous function $f : X \rightarrow [0, 1]$ such that $f(p) = 0$ and $f(B) = \{1\}$. Let G be a locally compact, Hausdorff, second countable group. A G -space is a topological space X with a given continuous action of G such that

- X is completely regular and any compact subset of X is metrizable.
- The quotient space X/G is paracompact and Hausdorff.

Note that the definition of a G -space in [1] uses the slightly more restrictive conditions that X and X/G be metrizable. Nothing is altered in [1] if this is replaced throughout with the above relaxed conditions.

The following definition may be found in Bourbaki [4, Definition 1, p.250, Prop.7, p.255].

Definition 3.1. The action of G on a G -space X is *proper* if given any two points $x, y \in X$ there are open neighbourhoods U_x, U_y of x and y respectively such that the set

$$\{g \in G : gU_x \cap U_y \neq \emptyset\}$$

has compact closure in G . A proper G -space X is said to be *G -compact* if the quotient X/G is compact.

In appendix A we give the full statement of Theorem 3.8 in Biller [3]. This theorem, on the existence of slices, reconciles the Bourbaki definition of proper actions with the definition in [1]. In particular, if X is a proper G -space such that X and X/G are metrizable then X satisfies the condition which was taken in [1] to be the definition of proper. The reverse implication is also valid. If X is a metrizable G -space which is proper in the sense of [1], then X is a proper G -space.

Note that if X is locally compact then this is equivalent to the following condition: if K_1, K_2 are compact subsets of X then the set

$$\{g \in G : gK_1 \cap K_2 \neq \emptyset\} \text{ is compact.}$$

It is easy to see any locally compact Hausdorff group acts properly on itself. Also note if X is a proper G -space and Y is any G -invariant subset of X then Y is a proper G -space when equipped with the subspace topology and obvious action of G . We shall want to define KK groups for the algebra $C_0(X)$ of a G -compact proper G -space and the following ensures that these algebras are separable.

Proposition 3.2. *Suppose G acts properly on X and X/G is compact. Then X is locally compact and second countable.*

Proof. The space X is locally compact by [4, (e), p.310]. We show that $X = GS$ for some compact $S \subset X$.

For each $x \in X$ let U_x be an open neighbourhood of x with compact closure. If $\pi : X \rightarrow X/G$ denotes the quotient map then the collection $\pi(U_x)$ cover X/G . We know X/G is compact and so get a finite sub-cover $\pi(U_{x_1}), \dots, \pi(U_{x_n})$. Now let $S = \cup_{i=1}^n \overline{U_{x_i}}$. This is a compact set with $GS = X$.

By assumption any compact subset of X is metrizable, and we may deduce that S is second countable when given the subspace topology from X .

The action of G on X gives rise to a map

$$G \times S \rightarrow GS = X,$$

which is continuous, open and surjective. By assumption G is second countable and so clearly X must be second countable. $\square$

Definition 3.3. Let X be a proper G -space. A *cutoff function* on X is a function $c : X \rightarrow \mathbb{R}_+$ such that the support of c has compact intersection with GK for any compact subset K of X and

$$\int c(g^{-1}x)dg = 1, \quad \text{for each } x \in X$$

Note the set of all such functions is convex.

Proposition 3.4. *Let X be a G -compact proper G -space. Then there exists a cutoff function on G .*

Proof. By Proposition 3.2 X is locally compact and $X = GS$ for some compact subset $S \subseteq X$. Take an open neighbourhood V of S with compact closure and let f be a continuous function from X into the interval $[0, 1]$ which is equal to 1 on S and zero on the complement of V , hence compactly supported.

Now for any $x \in X$ the function

$$f_x : G \rightarrow [0, 1], \quad f_x(g) = f(g^{-1}x)$$

is continuous. Because $GS = X$ we may find a $g \in G$ with $g^{-1}x \in S$. Recall $f = 1$ on S and so $f_x(g) = f(g^{-1}x) = 1$. As f_x is continuous we can find a neighbourhood of g on which f_x is non zero. Thus we may conclude

$$0 < \int f_x(g)dg$$

The action of G on X is proper so the set

$$\{g \in G : g\{x\} \cap \text{supp}(f) \neq \emptyset\}$$

is compact for each $x \in X$. This implies that f_x is compactly supported and so

$$\int f_x(g)dg < \infty.$$

Thanks to the above we may define

$$c(x) = \frac{f(x)}{\int f_x(g)dg}$$

This is compactly supported and has the property that $G \cdot \text{supp}(c) = X$ and

$$\int c(g^{-1}x)dg = 1$$

for each $x \in X$. Clearly c is a cutoff function on X . □

Definition 3.5. Let X, Y be proper G -spaces. A continuous map $\varphi : X \rightarrow Y$ is called a G -map if

$$\varphi(gx) = g\varphi(x) \text{ for all } g \in G, x \in X$$

Two G -maps are G -homotopic if they are homotopic through G -maps.

Definition 3.6. A universal example for proper actions of G , denoted $\underline{E}G$, is a proper G -space with the following property: If X is any proper G -space, then there exists a G -map $f : X \rightarrow \underline{E}G$, and any two such maps are G -homotopic.

Let K be any compact subset of G . The set of all probability measures on K — denoted $\mathcal{P}(K)$ — is a separable compact Hausdorff space in the topology induced from the weak* topology on $C(K)$. Recall $\mu_i \rightarrow \mu$ in the weak* topology if and only if

$$\int f d\mu_i \rightarrow \int f d\mu \text{ for any } f \in C(K).$$

If $K_1 \subseteq K_2$ are compact subgroups of G then define the following map

$$\iota : \mathcal{P}(K_1) \rightarrow \mathcal{P}(K_2), \quad (\iota\mu)(U) = \mu(U \cap K_1) \text{ for each Borel set } U \subseteq K_2$$

This is clearly injective and continuous. So ι is an injective map from a compact space to a Hausdorff space and hence gives rise to a homeomorphism between $\mathcal{P}(K_1)$ and its image in $\mathcal{P}(K_2)$ with the subspace topology. To simplify notation we identify $\mathcal{P}(K_1)$ with its image in $\mathcal{P}(K_2)$.

As G is locally compact and second countable it is clearly σ -compact and by [4, p.94, Prop. 15 and Cor. 2] we may find compact subsets

$K_1 \subset K_2 \subset K_3 \subset \dots$ with $G = \cup_{i=1}^{\infty} K_i$ such that *any* compact subset of G is contained within some K_i . Now define

$$\mathcal{P}_c(G) = \cup_{i=1}^{\infty} \mathcal{P}(K_i).$$

Clearly $\mathcal{P}_c(G)$ consists of all compactly supported probability measures on G . We topologize $\mathcal{P}_c(G)$ by giving it the direct limit topology with respect to the sequence $\mathcal{P}(K_i)$. This topology is independent of the sequence K_i .

The group G acts on $\mathcal{P}_c(G)$ by setting

$$(g\mu)(U) = \mu(g^{-1}U) \text{ for any Borel set } U \subseteq G.$$

Lemma 3.7. *The action of G on $\mathcal{P}_c(G)$ is continuous.*

Proof. Let $(g_\alpha, \mu_\alpha) \rightarrow (g, \mu)$, $\alpha \in A$, be a convergent net in $G \times \mathcal{P}_c(G)$. By definition of the direct limit topology on $\mathcal{P}_c(G)$, we must have $\mu_\alpha \rightarrow \mu$ in $\mathcal{P}(K_i)$ for some i . Given $f \in C(K_i)$ and $\epsilon > 0$ we can find α_0 with the property that

$$\alpha > \alpha_0 \Rightarrow \left| \int f d\mu_\alpha - \int f d\mu \right| \leq \epsilon/2$$

Furthermore $g_\alpha \rightarrow g$ and so we may choose α_0 in such a way that we also have

$$\alpha > \alpha_0 \Rightarrow \|f_{g_\alpha} - f_g\| \leq \epsilon/2$$

where f_g denotes the function $x \mapsto f(g^{-1}x)$. Finally note

$$\begin{aligned} \left| \int f d(g_\alpha \mu_\alpha) - \int f d(g\mu) \right| &= \left| \int f_{g_\alpha} d\mu_\alpha - \int f_g d\mu \right| \\ &\leq \left| \int (f_{g_\alpha} - f_g) d\mu_\alpha \right| + \left| \int f_g d\mu_\alpha - \int f_g d\mu \right| \\ &\leq \epsilon/2 + \epsilon/2 \end{aligned}$$

This shows that $g_\alpha \mu_\alpha \rightarrow g\mu$ in $\mathcal{P}(K_i)$ and therefore $g_\alpha \mu_\alpha \rightarrow g\mu$ in $\mathcal{P}_c(G)$. $\square$

Lemma 3.8. *The action of G on $\mathcal{P}_c(G)$ is proper.*

Proof. Take $\mu \in \mathcal{P}_c(G)$ and let f_μ be a continuous compactly supported function $f_\mu : G \rightarrow [0, 1]$ with $f_\mu \equiv 1$ on $\text{supp}(\mu)$.

Then the set

$$\begin{aligned} U_\mu &= \left\{ \lambda \in \mathcal{P}_c(G) : \left| \int f_\mu d\mu - \int f_\mu d\lambda \right| \leq 1/2 \right\} \\ &= \left\{ \lambda \in \mathcal{P}_c(G) : \int f_\mu d\lambda > 1/2 \right\} \end{aligned}$$

is an open neighbourhood of μ in $\mathcal{P}_c(G)$.

Now take any $\mu, \nu \in \mathcal{P}_c(G)$ and assume $gU_\mu \cap U_\nu \neq \emptyset$ for some $g \in G$. Indeed let $\lambda \in gU_\mu \cap U_\nu$. Then

$$\int g^{-1}f_\mu d\lambda > 1/2 \quad \text{and} \quad \int f_\nu d\lambda > 1/2$$

If $\text{supp}(g^{-1}f_\mu)$ and $\text{supp}(f_\nu)$ are disjoint then we have $0 \leq g^{-1}f_\mu + f_\nu \leq 1$. However by the above

$$\int g^{-1}f_\mu + f_\nu d\lambda > 1$$

clearly contradicting the fact that λ is a probability measure. Hence we may conclude that $g\text{supp}(\mu) \cap \text{supp}(\nu) \neq \emptyset$ and so

$$\{g \in G : gU_\mu \cap U_\nu \neq \emptyset\} \subset \{g \in G : g\text{supp}(\mu) \cap \text{supp}(\nu) \neq \emptyset\}.$$

Now both μ and ν are compactly supported and since any group acts properly on itself the larger set here is compact. $\square$

Theorem 3.9. *The space $\mathcal{P}_c(G)$ is a universal example for proper actions of G .*

Proof. Let X be any proper G -space, we aim to show there exists a G -equivariant map $X \rightarrow \mathcal{P}_c(G)$. Take any $x \in X$. By [3, Theorem 3.8] there is a G -invariant open neighbourhood U_x of x , a compact subgroup H of G , and a G -equivariant map

$$\rho : U_x \rightarrow G/H.$$

Let μ_H denote Haar measure on the compact subgroup H , normalized to have total mass 1. There is an obvious G -equivariant map

$$\Psi : G/H \rightarrow \mathcal{P}_c(G), \quad gH \mapsto g \cdot \mu_H$$

and let

$$\theta_x = \Psi \circ \rho : U_x \rightarrow \mathcal{P}_c(G).$$

Now X may be covered by such neighbourhoods and if π denotes the quotient map $X \rightarrow X/G$ then $\{\pi(U_x)\}$ is an open cover of X/G . Recall that by definition X/G is paracompact and Hausdorff and so there is a locally finite partition of unity subordinate to the cover $\{\pi(U_x)\}$. Precisely there are continuous maps

$$\omega_x : X/G \rightarrow [0, 1], \quad \text{for each } x \in X$$

with $\text{supp}(\omega_x) \subseteq \pi(U_x)$ and for each $[y] \in X/G$ we have

$$\omega_x([y]) = 0 \text{ for almost all } x \in X \text{ and } \sum_{x \in X} \omega_x([y]) = 1.$$

Now

$$\Xi : X \rightarrow \mathcal{P}_c(G), \quad \Xi(y) = \sum_{x \in X} \omega_x(\pi(y)) \theta_x(y)$$

is the required map. Note that (as remarked above) $\mathcal{P}_c(G)$ is a convex set and this convexity is being used in the construction of Ξ .

Finally, if $\varphi_1, \varphi_2 : X \rightarrow \mathcal{P}_c(G)$ are G -equivariant maps then they are G -homotopic via

$$t\varphi_1 + (1-t)\varphi_2, \quad t \in [0, 1].$$

So $\mathcal{P}_c(G)$ is a universal example for G . $\square$

Baum-Connes conjecture with coefficients. If A is any G - C^* -algebra then we may define

$$K_*^{\text{top}}(G, A) = \varinjlim_{\substack{G\text{-invariant} \\ G\text{-compact} \\ Z \subseteq \mathbb{E}G}} KK_*(C_0(Z), A)$$

We say a group G satisfies the Baum-Connes conjecture with coefficients if for every G - C^* -algebra A the map

$$\mu_A : K_*^{\text{top}}(G, A) \rightarrow K_*(A \rtimes_r G)$$

is an isomorphism.

4. K -THEORY FOR ASCENDING UNIONS OF GROUPS

Theorem 4.1. *Let H be an open subgroup of G . Then the inclusion of H in G determines a homomorphism of abelian groups*

$$\mathcal{T}_H^G : K_*(A \rtimes_r H) \rightarrow K_*(A \rtimes_r G).$$

Furthermore suppose $G_1 \subset G_2 \subset G_3 \subset \dots$ is an ascending sequence of open subgroups, then there is an inductive system of abelian groups

$$K_*(A \rtimes_r G_1) \xrightarrow{\mathcal{T}_{G_1}^{G_2}} K_*(A \rtimes_r G_2) \xrightarrow{\mathcal{T}_{G_2}^{G_3}} K_*(A \rtimes_r G_3) \xrightarrow{\mathcal{T}_{G_3}^{G_4}} \dots$$

If $G = \cup G_n$ we have

$$K_*(A \rtimes_r G) = \varinjlim_n K_*(A \rtimes_r G_n).$$

Proof. Any open subgroup H of G is also closed. Let H be an open subgroup of G and fix a Haar measure dg on G . Since H is an open subgroup of G , the restriction of dg to H is a Haar measure on H . To simplify notation we write dg to signify this choice of Haar measure on both G and H .

For $f \in C_c(H, A)$ define

$$\iota f : G \rightarrow A, \quad (\iota f)(x) = \begin{cases} f(x) & \text{if } x \in H \\ 0 & \text{otherwise.} \end{cases}$$

Then ι defines a map from $C_c(H, A)$ into $C_c(G, A)$ as H is clopen in G . ι is clearly additive and furthermore for $f, g \in C_c(H, A)$

$$\begin{aligned} \iota f * \iota g(x) &= \int_G \iota f(y) \alpha_y(\iota g(y^{-1}x)) dy \\ &= \int_H f(y) \alpha_y(\iota g(y^{-1}x)) dy \\ &= \begin{cases} \int_H f(y) \alpha_y(g(y^{-1}x)) dy & \text{if } x \in H \\ 0 & \text{otherwise.} \end{cases} \\ &= \iota(f * g)(x). \end{aligned}$$

Also for $f \in C_c(H, A)$

$$(\iota f)^*(x) = \begin{cases} \Delta_G(x^{-1}) \alpha_x(f(x^{-1})^*) & \text{if } x \in H \\ 0 & \text{otherwise} \end{cases} = (\iota f^*)(x)$$

This follows from the observation that $\Delta_H(x) = \Delta_G(x)$ for $x \in H$.

We now follow [6, II.C, p.172]. Let $\|\cdot\|_{A \rtimes_r G}$ signify the norm obtained on $C_c(G, A)$ from the left regular representation on the C^* -module

$$E = L^2(G, dg) \otimes A.$$

We shall also need the C^* -module

$$F = L^2(H, dg) \otimes A.$$

If $f \in C_c(H, A)$ then

$$\begin{aligned} \|\iota f\|_{A \rtimes_r G} &= \sup \{ \|\iota f * \xi\|_E : \xi \in E, \|\xi\|_E \leq 1 \} \\ &= \sup \{ \|f * \xi\|_F : \xi \in F, \|\xi\|_F \leq 1 \} \end{aligned}$$

So ι is an isometric $*$ -algebra map from $C_c(H, A)$ to $C_c(G, A)$. We can then complete this map to obtain an injective C^* -algebra morphism

$$\iota : A \rtimes_r H \rightarrow A \rtimes_r G.$$

The map $\mathcal{T}_H^G$ is then defined by functoriality:

$$\mathcal{T}_H^G = \iota_* : K_*(A \rtimes_r H) \rightarrow K_*(A \rtimes_r G).$$

Now let $G_1 \subset G_2 \subset G_3 \subset \dots$ be an open ascending sequence of groups and let $G = \cup G_n$. Fixing a Haar measure on G prescribes a Haar measure on each G_n . For any $m < n$, G_m is an open subgroup of G_n and so by the above we may define a map

$$\iota_m^n : A \rtimes_r G_m \rightarrow A \rtimes_r G_n.$$

Indeed we may form an inductive system of C^* -algebras $(A \rtimes G_n, \iota_m^n)$,

$$A \rtimes_r G_1 \xrightarrow{\iota_1^2} A \rtimes_r G_2 \xrightarrow{\iota_2^3} A \rtimes_r G_3 \xrightarrow{\iota_3^4} \dots$$

We aim to show that the C^* -inductive limit $\varinjlim_n A \rtimes_r G_n$ of this inductive system is equal to the algebra $A \rtimes_r G$. For any n , G_n is an open subgroup of G and let $\iota_n : A \rtimes_r G_n \rightarrow A \rtimes_r G$ be defined as above. Since each ι_n is injective, all that remains to be demonstrated is that

$$\bigcup_n \iota_n(A \rtimes_r G_n)$$

is dense in $A \rtimes_r G$.

Taking $\varepsilon > 0$ and $x \in A \rtimes_r G$ we can find $f \in C_c(G, A)$ with $\|f - x\|_{A \rtimes_r G} < \varepsilon$. The support of f is compact and by Lemma 2.1 must lie totally within some G_n . Hence $f \in C_c(G_n, A)$ for some n and so $\bigcup_n \iota_n(A \rtimes_r G_n)$ is dense in $A \rtimes_r G$. Hence $A \rtimes_r G$ is the C^* -inductive limit of the C^* -algebras $A \rtimes_r G_n$ and we therefore have

$$K_*(A \rtimes_r G) = \varinjlim_n K_*(A \rtimes_r G_n).$$

□

5. EQUIVARIANT K -HOMOLOGY FOR ASCENDING SEQUENCES OF GROUPS

Theorem 5.1. *Let H be an open subgroup of G . Then the inclusion of H in G determines a homomorphism of abelian groups*

$$\mathcal{R}_H^G : K_*^{\text{top}}(H, A) \rightarrow K_*^{\text{top}}(G, A).$$

Furthermore suppose $G_1 \subset G_2 \subset G_3 \subset \dots$ is an open ascending sequence of groups, then there is an inductive system of abelian groups

$$K_*^{\text{top}}(G_1, A) \xrightarrow{\mathcal{R}_{G_1}^{G_2}} K_*^{\text{top}}(G_2, A) \xrightarrow{\mathcal{R}_{G_2}^{G_3}} K_*^{\text{top}}(G_3, A) \xrightarrow{\mathcal{R}_{G_3}^{G_4}} \dots$$

If $G = \bigcup G_n$ then we have

$$K_*^{\text{top}}(G, A) = \varinjlim_n K_*^{\text{top}}(G_n, A)$$

In the course of proving this result, we shall make use of the *reciprocity isomorphism* [5, p.157] in equivariant KK -theory. Let H be an open subgroup of G , let A be an H - C^* -algebra, let $\text{Ind}_H^G A$ denote the induced algebra and let B be a G - C^* -algebra. Then we have the reciprocity isomorphism:

$$\text{Inf}_H^G : KK_*^H(A, B) \cong KK_*^G(\text{Ind}_H^G A, B).$$

If A is commutative we have $A \cong C_0(X)$, $\text{Ind}_H^G A \cong C_0(G \times_H X)$ and so we have

$$\text{Inf}_X : K_*^H(C_0(X), B) \rightarrow K_*^G(C_0(G \times_H X), B).$$

Lemma 5.2. *Let H be an open subgroup of G and let X be any H -compact subset of $\mathcal{P}_c(H) \subset \mathcal{P}_c(G)$ then $G \times_H X \cong G \cdot X$ as G -spaces.*

Proof. Recall the definition of the space $G \times_H X$. The group H acts on the product $G \times X$ by setting $h \cdot (g, x) = (gh^{-1}, hx)$ and $G \times_H X$ is the quotient. The action of G on $G \times_H X$ is given by $g' \cdot [g, x] = [g'g, x]$, where $[g, x]$ is the equivalence class of the pair (g, x) .

The map F_X is defined as follows:

$$F_X : G \times_H X \rightarrow G \cdot X, [g, x] \mapsto gx.$$

The map F_X is clearly surjective and G -equivariant. To show this map is injective take $[g_1, x_1], [g_2, x_2] \in G \times_H X$ with $g_1 x_1 = g_2 x_2$. Recall that x_1, x_2 are in fact probability measures on H and so

$$1 = x_1(H) = g_1^{-1} g_2 x_2(H) = x_2(g_2^{-1} g_1 H)$$

As x_2 is a measure in $\mathcal{P}_c(H)$ it will clearly be equal to zero on the coset $g_2^{-1} g_1 H$ unless $g_2^{-1} g_1 \in H$. Now $g_2^{-1} g_1 \cdot (g_1, x_1) = (g_1 g_1^{-1} g_2, g_2^{-1} g_1 x_1) = (g_2, x_2)$ as required.

It now remains to show this map is a homeomorphism. Let π denote the quotient map $G \times X \rightarrow G \times_H X$ and $\theta : G \times X \rightarrow G \cdot X$ denote the map given by the action of G on X . Both π and θ are open, continuous maps. Since $\theta = F_X \circ \pi$, this implies that F_X is open and continuous. $\square$

Let $\mathcal{R}_{H,X}^G$ denote the composition

$$KK_*^H(C_0(X), A) \xrightarrow{\text{Inf}_X} KK_*^G(C_0(G \times_H X), A) \xrightarrow{(F_X)_*} KK_*^G(C_0(G \cdot X), A)$$

If X and Y are H -compact subsets of $\mathcal{P}_c(H)$ with $X \subset Y$ then the following diagram commutes,

$$\begin{array}{ccc} KK_*^H(C_0(X), A) & \longrightarrow & KK_*^H(C_0(Y), A) \\ \mathcal{R}_{H,X}^G \downarrow & & \mathcal{R}_{H,Y}^G \downarrow \\ KK_*^G(C_0(G \cdot X), A) & \longrightarrow & KK_*^G(C_0(G \cdot Y), A) \end{array}$$

where each horizontal map is given by inclusion of the spaces involved.

Now $\mathcal{P}_c(H), \mathcal{P}_c(G)$ are universal examples for H, G respectively. If X is an H -compact subset of $\mathcal{P}_c(H)$ then $G \cdot X$ is a G -compact subset of $\mathcal{P}_c(G)$. Due to the above the following map is well defined

$$\mathcal{R}_H^G : K_*^{\text{top}}(H, A) \rightarrow K_*^{\text{top}}(G, A).$$

Now let $G_1 \subset G_2 \subset G_3 \subset \dots$ be an ascending sequence of open subgroups and let $G = \cup G_n$. There is then an inductive system of abelian groups

$$K_*^{\text{top}}(G_1, A) \xrightarrow{\mathcal{R}_{G_1}^{G_2}} K_*^{\text{top}}(G_2, A) \xrightarrow{\mathcal{R}_{G_2}^{G_3}} K_*^{\text{top}}(G_3, A) \xrightarrow{\mathcal{R}_{G_3}^{G_4}} \dots$$

and the maps

$$\mathcal{R}_{G_n}^G : K_*^{\text{top}}(G_n, A) \rightarrow K_*^{\text{top}}(G, A).$$

Lemma 5.3. *There exist compact sets $\Delta_1 \subset \Delta_2 \subset \Delta_3 \subset \dots$ such that $\Delta_n \subset G_n$ and $\cup_n \text{Interior}(\Delta_n) = G$. Set $Z^{n,m} = G_n \cdot P_c(G_n \cap \Delta_m)$. Then*

- $Z^{n,m} \subset P_c(G_n)$
- $Z^{n,m}$ is preserved by G_n and is G_n -compact
- $Z^{n,m} \subset Z^{n,m+1}$
- $\cup_m Z^{n,m} = P_c(G_n)$
- $G_{n+1} \cdot Z^{n,m} \subset Z^{n+1,m}$

Proof. Let $W_1, W_2, W_3, \dots$ be a countable basis for the topology of G . From this list $W_1, W_2, W_3, \dots$ delete all W_j such that $\overline{W_j}$ is not compact. Call the remaining list $V_1, V_2, V_3, \dots$. Let U be any non-empty open set in G and choose $p \in U$. Since G is locally compact there exists an open set Λ in Y with $p \in \Lambda$ and $\overline{\Lambda}$ compact. Consider $\Lambda \cap U$. Now $\Lambda \cap U$ is a union of W_j 's. But any W_j which is contained in $\Lambda \cap U$ must in fact be a V_k because $\Lambda \cap U$ has compact closure. $\Lambda \cap U$ is a union of V_k 's (now let p vary throughout U).

Now let $\Sigma_n = \overline{V_1 \cup V_2 \cup \dots \cup V_n}$ and $\Delta_n = G_n \cap \Sigma_n$.

Any compact set in G is contained in some Δ_n . □

Lemma 5.4. *Let $\{A^{m,n}\}$ be a commutative diagram of abelian groups in which the typical commutative square is*

$$\begin{array}{ccc} A^{n,m} & \longrightarrow & A^{n,m+1} \\ \downarrow & & \downarrow \\ A^{n+1,m} & \longrightarrow & A^{n+1,m+1} \end{array}$$

with $n, m = 1, 2, 3, \dots$. Then there is a canonical isomorphism of abelian groups:

$$\lim_{m \rightarrow \infty} (\lim_{n \rightarrow \infty} A^{m,n}) \cong \lim_{n \rightarrow \infty} (\lim_{m \rightarrow \infty} A^{m,n}).$$

Proof. Each side is canonically isomorphic to the direct limit $\varinjlim A^{n,n}$ of the directed system $\{A^{n,n}\}$. $\square$

Theorem 5.5. $\lim_{n \rightarrow \infty} K_*^{top}(G_n, A) = K_*^{top}(G, A)$.

Proof. Consider the commutative diagram of abelian groups in which the typical commutative square is:

$$\begin{array}{ccc} \mathcal{K}_j^{G_n}(Z^{n,m}, A) & \longrightarrow & K_j^{G_n}(Z^{n,m+1}, A) \\ \rho_n \downarrow & & \rho_n \downarrow \\ K_j^{G_{n+1}}(Z^{n+1,m}, A) & \longrightarrow & K_j^{G_{n+1}}(Z^{n+1,m+1}, A) \end{array}$$

with $n, m = 1, 2, 3, \dots$. Each horizontal arrow

$$K_j^{G_n}(Z^{n,m}, A) \rightarrow K_j^{G_n}(Z^{n,m+1}, A)$$

is the map of abelian groups determined by the inclusion $Z^{n,m} \rightarrow Z^{n,m+1}$. Each vertical map $\rho_n : K_j^{G_n}(Z^{n,m}, A) \rightarrow K_j^{G_{n+1}}(Z^{n+1,m}, A)$ is the map $\mathcal{R}_{G_n}^{G_{n+1}}$ followed by the map of abelian groups induced by the inclusion $G_{n+1} \cdot Z^{n,m} \rightarrow Z^{n+1,m}$.

We will write

$$A^{n,m} = K_j^{G_n}(Z^{n,m}, A).$$

Each $Z^{n,m}$ is G_n -compact, $\cup_m Z^{n,m} = P_c(G_n)$ and any G_n -compact set in $P_c(G_n)$ is contained in some $Z^{n,m}$. Taking the direct limit along the n th row, we have

$$\lim_{m \rightarrow \infty} A^{m,n} = K_j^{top}(G_n, A).$$

If we now take the direct limit in a vertically downward direction, we obtain

$$\lim_{n \rightarrow \infty} K_j^{top}(G_n, A).$$

Now we fix attention on the m th column. If $n \geq m$ then $G_n \supset \Delta_m$ and so $G_n \cap \Delta_m = \Delta_m$. Then we have

$$A^{n,m} = K_j^{G_n}(G_n \cdot P_c(\Delta_m), A).$$

Now the G_n -saturation of $P_c(\Delta_m)$ is equal to the G -saturation of $P_c(\Delta_m)$. Now we apply the reciprocity isomorphism and we have

$$A^{n,m} \cong K_j^G(G \cdot P_c(\Delta_m), A).$$

if $n \geq m$. Therefore the m th column *stabilizes* as soon as $n \geq m$. Therefore the direct limit down the m th column is given by

$$\lim_{n \rightarrow \infty} A^{m,n} = K_j^G(G \cdot P_c(\Delta_m), A).$$

Now the sets $G \cdot P_c(\Delta_m)$ are cofinal in G -compact sets in $P_c(G)$. Taking the direct limit in the horizontal direction, we have

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} A^{m,n} \cong K_j^{\text{top}}(G, A).$$

By Lemma 5.4 we have

$$\lim_{n \rightarrow \infty} K_j^{\text{top}}(G_n, A) \cong K_j^{\text{top}}(G, A).$$

□

6. ADELIC GROUPS

For H an open subgroup of G we have constructed a homomorphism from $K_*^{\text{top}}(H, A)$ to $K_*^{\text{top}}(G, A)$, and likewise for the K theory of the reduced crossed product C^* -algebras. We wish to check that these maps are compatible with the Baum–Connes μ map, i. e. that the following diagram commutes.

$$\begin{array}{ccc} K_*^{\text{top}}(H, A) & \xrightarrow{\mu_H} & K_*(A \rtimes_r H) \\ \mathcal{R}_H^G \downarrow & & \downarrow \mathcal{T}_H^G \\ K_*^{\text{top}}(G, A) & \xrightarrow{\mu_G} & K_*(A \rtimes_r G) \end{array}$$

As a first step, we prove that the reciprocity isomorphism Inf_X is compatible with the Baum–Connes μ map.

Lemma 6.1. *Let A be a $G - C^*$ -algebra, let H be an open subgroup of G and let X be a locally compact proper H -compact H -space. Then the following diagram commutes:*

$$\begin{array}{ccc} KK_*^H(C_0(X), A) & \xrightarrow{\mu_H} & K_*(A \rtimes_r H) \\ \text{Inf}_X \downarrow & & \downarrow \mathcal{T}_H^G \\ KK_*^G(C_0(G \times_H X), A) & \xrightarrow{\mu_G} & K_*(A \rtimes_r G) \end{array}$$

Proof. The inverse of the reciprocity isomorphism Inf is the *compression isomorphism* [5, p. 157] and is given by the composition $i_* \circ \text{Res}_X$, where Res_X is the obvious restriction map $KK_*^G(C_0(X), A) \rightarrow KK_*^H(C_0(X), A)$ and i is the inclusion $C_0(X) \hookrightarrow C_0(G \times_H X)$ given by

$$i(f)[g, x] = \begin{cases} f(gx) & \text{if } g \in H \\ 0 & \text{otherwise.} \end{cases}$$

Each of these maps is clearly functorial.

There is a commutative diagram

$$\begin{array}{ccc}
 KK_*^H(C_0(X), A) & \xrightarrow{j_H} & KK_*(C_0(X) \rtimes_r H, A \rtimes_r H) \\
 i_* \uparrow & & i'_* \uparrow \\
 KK_*^H(C_0(G \times_H X), A) & \xrightarrow{j_H} & KK_*(C_0(G \times_H X) \rtimes_r H, A \rtimes_r H) \\
 \text{Res}_X \uparrow & & p_* \circ q_* \downarrow \\
 KK_*^G(C_0(G \times_H X), A) & \xrightarrow{j_G} & KK_*(C_0(G \times_H X) \rtimes_r G, A \rtimes_r G)
 \end{array}$$

in which j_H, j_G are *descent* homomorphisms. Here p denotes the map

$$p : C_0(G \times_H X) \rtimes_r G \rightarrow C_0(G \times_H X) \rtimes_r H$$

induced from the obvious restriction map $C_c(G, A) \rightarrow C_c(H, A)$, with $A = C_0(G \times_H X)$. And q denotes the map

$$q : A \rtimes_r H \rightarrow A \rtimes_r G$$

of Theorem 4.1.

Now let c be a cutoff function on the proper H -space X , we claim $i(c) \in C_0(G \times_H X)$ is a cutoff function on the proper G -space $G \times_H X$. To see this take any $[g_0, x] \in G \times_H X$. Then by definition we have

$$i(c)([g_0, x]) = 0$$

unless $g_0 \in H$ in which case

$$i(c)[g_0, x] = c(g_0 x).$$

Up to a normalizing factor between the Haar measures on H and G

$$\begin{aligned}
 \int_G i(c)(g^{-1}[g_0, x]) \, dg &= \int_G i(c)(g^{-1}g_0^{-1}[g_0, x]) \, dg \\
 &= \int_G i(c)([g^{-1}, x]) \, dg \\
 &= \int_H c(h^{-1}x) \, dh = 1.
 \end{aligned}$$

If K is any compact subset of $G \times_H X$ and if F denotes the homeomorphism between $G \times_H X$ and $G \cdot X$ then $F(GK) = G \cdot F(K)$ and $F(K)$ is compact. Also note $F(\text{supp}(i(c))) \subseteq H \cdot \text{supp}(c)$ and so

$$F(GK \cap \text{supp}(i(c))) \subseteq G \cdot F(K) \cap H \cdot \text{supp}(c) \subseteq H \cdot F(K) \cap \text{supp}(c)$$

which is compact and so $GK \cap \text{supp}(i(c))$ is compact. So we have shown $i(c)$ is a cutoff function on $G \times_H X$.

Let λ_X denote the projection in the twisted convolution algebra $C_c(H \times X)$ arising from the cutoff function c :

$$\lambda_Z(g, x) = c_Z(x)^{1/2} c_Z(g^{-1}x)^{1/2} \Delta(g)^{-1/2}$$

and let $\lambda_{G \times_H X}$ denote the projection in $C_c(G \times G \times_H X)$ arising from the cutoff function $i(c)$.

Then $p(\lambda_{G \times_H X})$ is simply the restriction of $\lambda_{G \times_H X}$ to $H \times G \times_H X$, and for any h in G and any $[g, x] \in G \times_H X$

$$\begin{aligned} p(\lambda_{G \times_H X})(h, [g, x]) &= i(c)[g, x]^{1/2} i(c)(h^{-1}[g, x])^{1/2} \Delta_G(h)^{-1/2} \\ &= \begin{cases} c(gx)^{1/2} c(h^{-1}gx) \Delta_H(h)^{-1/2} & \text{if } g \in H \\ 0 & \text{otherwise.} \end{cases} \\ &= i'(\lambda_X) \end{aligned}$$

So the following diagram commutes

$$\begin{array}{ccc} KK_*(C_0(X) \rtimes_r H, A \rtimes_r H) & \xrightarrow{[\lambda_X] \otimes \cdot} & KK_*(\mathbb{C}, A \rtimes_r H) \\ i'_* \uparrow & & \parallel \\ KK_*(C_0(G \times_H X) \rtimes_r H, A \rtimes_r H) & \xrightarrow{i'_*([\lambda_X]) \otimes \cdot} & KK_*(\mathbb{C}, A \rtimes_r H) \\ p_* \circ q_* \downarrow & & q_* \downarrow \\ KK_*(C_0(G \times_H X) \rtimes_r G, A \rtimes_r G) & \xrightarrow{[\lambda_{G \times_H X}] \otimes \cdot} & KK_*(\mathbb{C}, A \rtimes_r G) \end{array}$$

We finish the proof by splicing together these two diagrams. $\square$

Lemma 6.2. *Let X be an H -compact subset of $P_c(H)$. Then the following diagram commutes:*

$$\begin{array}{ccc} KK_*^H(C_0(X), A) & \xrightarrow{\mu_H} & K_*(A \rtimes_r H) \\ \mathcal{R}_{H,X}^G \downarrow & & \mathcal{T}_H^G \downarrow \\ KK_*^G(C_0(G \cdot X), A) & \xrightarrow{\mu_G} & K_*(A \rtimes_r G). \end{array}$$

Proof. By Lemma 5.2 we know that the induced space $G \times_H X$ is G -homeomorphic to the G -saturation $G \cdot X$. We now apply Lemma 6.1. $\square$

Theorem 6.3. *Let G be a locally compact, second countable Hausdorff topological group and let A be a G - C^* -algebra. Let G be the union of open subgroups G_n such that the Baum-Connes conjecture with coefficients A is true for each G_n . Then the Baum-Connes conjecture with coefficients A is true for G .*

Proof. We start with the commutative diagram in Lemma 6.2 and take the direct limit over all H -compact subsets of $P_c(H)$. We then obtain the commutative diagram

$$\begin{array}{ccc} K_*^{\text{top}}(H, A) & \xrightarrow{\mu_H} & K_*(A \rtimes_r H) \\ \mathcal{R}_H^G \downarrow & & \downarrow \mathcal{T}_H^G \\ K_*^{\text{top}}(G, A) & \xrightarrow{\mu_G} & K_*(A \rtimes_r G) \end{array}$$

If G is the union of open subgroups G_i each of which satisfies *BCC*, then applying Theorem 4.1 and Theorem 5.1 along with the above commutative diagram is enough to show that G satisfies *BCC*. $\square$

Theorem 6.4. *Let F be a global field, $\mathbb{A}$ its ring of adeles, G a linear algebraic group over F . Let F_v denote a place of F . If *BCC* is true for each local group $G(F_v)$ then *BCC* is true for the adelic group $G(\mathbb{A})$.*

Proof. Let $v_1, v_2, v_3, \dots$ be an ordering of the finite places of F , let S_∞ denote the finite set of all infinite places of F , and let $S(n) = \{v_1, v_2, \dots, v_n\} \cup S_\infty$. Let $G_n = G_{S(n)}$ in the notation of section 2.

If Γ is a compact group then $\underline{E}\Gamma$ is a point, and we have

$$K_*^{\text{top}}(\Gamma, B) \cong KK_*^\Gamma(B) \cong K_*(B \rtimes \Gamma)$$

by the Green-Julg theorem. Therefore *BCC* is true for any compact group. Now G_n is a product of finitely many local groups and one compact group. But *BCC* is stable under the direct product of finitely many groups, by an important result of Chabert-Echterhoff [5, Theorem 3.17]. Therefore *BCC* is true for each *open* subgroup G_n . Now the adelic group $G(\mathbb{A})$ is the ascending union of the open subgroups G_n , therefore *BCC* is true for $G(\mathbb{A})$, by Theorem 6.3. $\square$

APPENDIX A. BILLER'S THEOREM

We give here the full statement of Theorem 3.8 in Biller[3]. The stabilizer of $x \in X$ is denoted G_x .

Theorem (Existence of slices). Let G be a locally compact group acting properly on a completely regular space X , and choose $x \in X$. Then there is a convergent filter basis $\mathcal{N}$ that consists of compact subgroups of G normalized by G_x such that for every $N \in \mathcal{N}$, the coset space $G/G_x N$ is a manifold and x is contained in a $G_x N$ -slice for the action of G on X . In particular, some neighbourhood of the orbit $G \cdot x$ is a locally trivial fibre bundle over the manifold $G/G_x N$.

The dimension of $G \cdot x$ is infinite if and only if $N \in \mathcal{N}$ may be chosen such that the dimension of $G/G_x N$ is arbitrarily high. If the

dimension of $G \cdot x$ is finite, then $N \in \mathcal{N}$ may be chosen in such a way that $\dim G/G_x N = \dim G \cdot x$.

REFERENCES

- [1] P. Baum, A. Connes, and N. Higson. Classifying space for proper actions and K -theory of group C^* algebras. *Contemporary Math.* 167 (1994) 241–291.
- [2] P. Baum, S. Millington, and R. Plymen. A proof of the Baum-Connes conjecture for reductive adelic groups. *C. R. Math. Acad. Sci. Paris*, 332 (2001) 195–200.
- [3] H. Biller, Proper actions on cohomology manifolds, *Trans. Amer. Math. Soc.* 355 (2003) 407 – 432.
- [4] N. Bourbaki. *General Topology*. Addison-Wesley, 1966.
- [5] J. Chabert and S. Echterhoff. Permanence properties of the Baum-Connes conjecture. *Doc. Math.* 6 (2001) 127–183.
- [6] A. Connes. *Noncommutative geometry*. Academic Press, New York, 1994.
- [7] J. Elstrodt, F. Grunewald, J. Mennicke, Groups acting on hyperbolic space, Springer, Berlin 1998.
- [8] N. Higson and G. G. Kasparov. E -theory and KK -theory for groups which act properly and isometrically on Hilbert space. *Invent. Math.* 144 (2001) 23–74
- [9] P. Julg, Travaux de N. Higson et G. Kasparov sur la conjecture de Baum-Connes, *Astérisque* 252 (1998) 151 –183.
- [10] P. Julg, The Baum-Connes conjecture with coefficients for the group $Sp(n, 1)$, *C. R. Math. Acad. Sci. Paris* 334 (2002) 533 –538.
- [11] P. Julg and G.G. Kasparov, Operator K -theory for the group $SU(n, 1)$, *J. Reine Angew. Math.* 463 (1995) 99 –152.
- [12] G.G. Kasparov, Lorentz groups: K -theory of unitary representations and crossed products, *Soviet Math. Dokl.* 27 (1983) 105 – 109.
- [13] J-P. Serre, *Trees*, Springer, Berlin 1980.
- [14] A. Weil. *Adeles and algebraic groups*. Birkhauser, 1982.
- [15] A. Weil, *Basic Number Theory*, Springer Classics in Math. 1995.

Paul Baum, Mathematics Department, Pennsylvania State University, University Park, PA 16802, USA. Email: baum@math.psu.edu

Stephen Millington, Compass Computer Consultants Ltd, Compass House, Frodsham, Cheshire WA6 0AQ, UK. Email: stem@quick.freemove.co.uk

Roger Plymen, Mathematics Department, Manchester University, Manchester M13 9PL, UK. Email: roger@ma.man.ac.uk